Clinical outcome of Descemet membrane endothelial keratoplasty (DMEK) with imported donor corneas in eyes of Asian patients; endothelium-in versus endothelium-out method

Young-ho Jung, Chang Ho Yoon, Mee Kum Kim

Published: June 30, 2022 • https://doi.org/10.1371/journal.pone.0270037

Abstract

Objective

We investigated whether (1) imported pre-cut tissue is feasible for Descemet membrane endothelial keratoplasty (DMEK) in eyes of Asian patients, (2) the clinical outcome is comparable between the endothelium-in and endothelium-out methods, and (3) the corneal edema-induced anterior curvature changes may have an effect on the refractive error.

Methods

The medical records of 32 DMEK patients who underwent either the endothelium-out or endothelium-in method using imported precut grafts with a 3-day pre-cut-to-use time were retrospectively analyzed. Fuchs' endothelial dystrophy (37.5%) and bullous keratopathy (62.5%) cases were included. The main clinical outcome measures were graft survival, best corrected visual acuity (BCVA), endothelial cell density (ECD), corneal thickness (CT), and complications. Correlation of the anterior curvature changes with refractive error was analyzed in the DMEK with cataract surgery group.

Results

The overall survival rate was 71.9%. Final graft failures were caused by rejection, glaucoma, and infection. Visual acuities improved by 89.3%. BCVA better than 20/40 and 20/20 was found in 75% and 28.6% of patients, respectively. The ECDs at 3 months and 1 year were 1400 and 1083 cells/mm², respectively. The mean survival time, ECD, BCVA, CT, and complication rates were not different between the endothelium-in and endothelium-out methods. A hyperopic shift by +0.42 D was not related to the anterior curvature changes.

Conclusion

Imported pre-cut tissues with a ≤ 3-day pre-cut-to-use time are feasible for DMEK in the treatment of corneal endothelial edema in eyes of Asian patients, and both endothelium-in and endothelium-out methods appear to be comparatively effective. Edema-induced anterior curvature change may not affect the refractive shift.

Citation: Jung Y-h, Yoon CH, Kim MK (2022) Clinical outcome of Descemet membrane endothelial keratoplasty (DMEK) with imported donor corneas in eyes of Asian patients; endothelium-in versus endothelium-out method. PLoS ONE 17(6): e0270037. https://doi.org/10.1371/journal.pone.0270037

Editor: Rajiv R. Mohan, University of Missouri-Columbia, UNITED STATES

Received: September 9, 2021; Accepted: June 2, 2022; Published: June 30, 2022

Copyright: © 2022 Jung et al. This is an open access article distributed under the terms of the <u>Creative Commons Attribution</u> <u>License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: All relevant data are within the paper and its Supporting Information files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Since Gerrit Melles first proposed the concept of posterior lamellar keratoplasty in 1998 [1], endothelial keratoplasty has been prevalent in Western countries for both bullous keratopathy (BK) and Fuchs' endothelial dystrophy (FED) [2, 3]. After the introduction of Descemet membrane endothelial keratoplasty (DMEK) by Gerrit Melles [4-6], faster and better visual recovery and a lower rate of graft rejection were noted, compared to either Descemet stripping automated endothelial keratoplasty (DSAEK) or penetrating keratoplasty [2, 3, 7]. Currently, nearly 850 articles related to DMEK are available on PubMed. Among them, hundreds of articles present the clinical outcomes of DMEK in Western countries. There are presently only about 20 articles that describe the clinical outcomes of DMEK in Asian countries. The low popularity of DMEK in Asia could be related to multiple causes such as the low surgical volume owing to the shortage of donor corneas, the surgically challenging small dimension of the eyes, and the rarity of FED cases in Asia [8-13].

Since various surgical techniques are employed in DMEK to reduce the endothelial cell loss [8, 14–19], standardization of the DMEK procedure is necessary, especially for the challenging cases of Asian eyes. In Korea, there are very few native corneal donors. Therefore, patients who wish to undergo DMEK prefer opting for imported tissues to ensure a timely operation rather than waiting longer for a native tissue. Unlike Western countries, where pre-cut tissue is available within 24–48 hours following the stripping procedure, overseas transport of pre-cut tissues from eye banks in Western countries to Asia takes more than 2 days. It is not fully elucidated whether the overseas transport of pre-cut tissues, with more than a 48-hour period between the pre-cutting and the use time, affects the clinical outcome of DMEK in eyes of Asian patients.

Additionally, the refractive outcome and factors causing shift in the refractive errors have been currently debated in patients with FED following DMEK combined with cataract surgery [20–23]. Various incidents of hyperopic surprise have been reported, and preoperative thicker cornea or postoperative flatter posterior curvature have been suspected to be the underlying risk factors [20, 22]. However, it is still not clear whether edema-induced anterior curvature changes have an effect on the refractive errors in patients with corneal edema.

Overall, three issues need to be addressed: (1) whether imported pre-cut tissues are feasible for use in DMEK in Asia, (2) whether the clinical efficacy of the endothelium-in delivery method is comparable to that of the endothelium-out method in DMEK, and (3) the extent of shift in the refractive error following DMEK and whether it is related to the anterior curvature changes. Therefore, this study investigated 1) the clinical efficacy of DMEK in eyes of Asian patients using imported pre-cut tissues with a \leq 3-day pre-cut-to-use time, 2) the correlation of the anterior curvature changes with refractive shift in DMEK combined with cataract surgery, and 3) whether the clinical efficacy of the endothelium-in delivery is comparable to that of the endothelium-out method in DMEK with imported pre-cut tissues. These results will benefit the standardization of DMEK procedures in eyes of Asian patients using imported pre-cut tissues.

Materials and method

Subjects

The study adhered to the tenets of the Declaration of Helsinki. Informed consent was waived by the Institutional Review Board because the study was based on the retrospective review of old charts. Under the approval by the Institutional Review Board of Seoul National University Hospital (IRB No. 2108-026-1241), the medical records of DMEK cases performed by one surgeon (M.K.K) at our tertiary referral center between October in 2016 and May 6th in 2022 were retrospectively reviewed.

Patients who had been diagnosed with corneal endothelial edema due to either FED or BK and had undergone DMEK with ≥ 6 month-follow up were included. The surgical indication was persistent corneal stromal edema with decreased visual acuity. Among the 35 DMEK cases in the above-mentioned period, three cases were excluded from the analysis. Two were excluded owing to loss in follow-up (< 6 months), and the remaining was excluded because the patient had previously undergone whole corneal transplantation prior to DMEK with a high risk of rejection. Of the total 32 cases, one-third had grade 4 FED with corneal stromal edema (n = 12, 37.5%), and two-third had BK (n = 20, 62.5%) [24, 25]. Eyes with previous posterior capsular openings or those that had undergone a previous vitrectomy were also included. (n = 4, 12.5%). DMEK combined with cataract surgery was performed in 18 cases, concomitantly (2 cases) or sequentially (16 cases). Pupilloplasty was performed sequentially in 3 cases before DMEK. All the patients' demographics are presented in Table 1. Although the indications of the patients were heterogenous (either FED or bullous keratopathy) in each group, there was no statistical difference in the distribution of the previous diagnosis between these two groups.

Characteristics	Total (n = 32)	Ende-out (n = 17)	Endo in (n = 15)	P value
Recipients				
Apr (years)	64.5x11.1	62.9110.8	46.3x11.5	0.396
Ses (MDF)	13:29	6.11	7:8	6.720
Prosperative BCVA (logMAR)	0.5210.4	0.94±0.4	1.0110.4	8.630
Surgical indication (Fuchs' vs BK)	12:30	7,10	5: 10	8.647
Combined surgery				
Cataract surgery	18 (56.3)	11 ()(4.7)	7 (467)	8.621
Pupilloplasty	3 (9.4)	2(11.8)	1 (6.7)	0.476
Ocular comorbidities	28 (42.5)	10 (58.8)	18 (96.7)	8.647
Glaucoma (n, %)	11 (14.4)	7(41.2)	4 (26.7)	0.388
Other comorbidities	11 (34.4)	4 (29.5)	7 (46.7)	0.149
ACD (phakic state)	2.2e8.6	2.2+0.7	2.1ell-6	0.748
AXL (mm)	24.1±3.2	244±3.5	23.3 ± 2.5	0.518
CCT (µm)	677163.1	6764.99.8	487168.8	8.905
ECD (sells/mm ²)*	484±80.3	486+94.7	498±13.4	4.903
Follow-up (months)	33.3±15.7	42.3±11.8	23.6x7.1	< 0.001
Nations				
Apr (years)	56.2±4.6	57.8±4.1	54304.6	0.043
Graft size (mm)	7.72x0.5	7.4x8.3	8.010.4	<0.001
Donor ECD (cells/mm ²)	2812+205	2837+169	2783+243	0.469
Death-to-use time (hours)	121111	127x12	1354.8	0.051
Pre-cut to use time	59+15	59+15	58+15	0.951

Ends endelschem 34, mehr 15, fendels RCVA, bei cernord visual early BK, ballas konzupelly, KCX, anders chandre depth, AX, and i angeh, ECX, anderheit of density, OHm methics include events is andere segarest synthese, epitential mechanise, systaat mechanise, and pendendifiation syntheses etc. BCIP role susceed for score 2029 avoing to cortexel eleman.

Table 1. Preoperative demographics in recipients and donors. https://doi.org/10.1371/journal.pone.0270037.t001

The following data were collected from the medical charts: demographic information that included the recipient's age, sex, general medical history, ocular medical and surgical history; and ophthalmic findings that included visual acuity, intraocular pressure (IOP), endothelial cell density (ECD), central corneal thickness (CCT), ocular biometric parameters, and the follow-up period (<u>Table 1</u>). Concomitant ophthalmic diseases that might affect graft survival or visual acuity, which included glaucoma, uveitis, toxic anterior segment syndrome, epiretinal membranes, age-related macular degeneration, cystoid macular edema, pseudoexfoliation syndrome, pupillary abnormality, and iris synechiae are shown in <u>Fig 1</u>.



Fig 1. Representative photographs of patients with a concomitant ocular disease who underwent Descemet membrane endothelial keratoplasty (DMEK).

First and second column: preoperative photographs of the cornea. Third column: horizontal (A–E) or vertical scans (F) of anterior segment optical coherence tomography. Fourth column: postoperative photographs of the cornea after DMEK. (A) Anterior displaced crystalline lens due to zonular weakness. (B) Posterior iris synechiae (red arrows) after acute angle-closure glaucoma. (C) Dilated and paralyzed pupil (white arrowhead), keratic precipitates (red arrows) and shallow anterior chamber due to acute angle-closure glaucoma. Pupilloplasty and cataract surgery were performed prior to DMEK. (D) Iris-fixed phakic intraocular lens (IOL)-induced iritis and bullous keratopathy. (E) Uveitis glaucoma with thickened corneal endothelium (red arrows) and peripheral anterior synechiae (white arrows). Glaucoma drainage device insertion (white arrowhead) after DMEK. (F) Toxic anterior segment syndrome and uvietic glaucoma with progressive peripheral anterior synechiae (white arrows). Dilated pupil (white arrowhead) is corrected with pupilloplasty (red arrow). https://doi.org/10.1371/journal.pone.0270037.g001

Donor characteristics in overseas transport of pre-cut tissues

A total of 32 transplant grafts were supplied by the Eversight Eye Bank (Chicago, IL) using the standard procedure [<u>26</u>, <u>27</u>]. Only donor tissues with a death-to-use period of < 7 days were imported. Double partial trephinated-grafts with S-marks were used. The size of the grafts ranged from 7.25–8.5 mm and the grafts were selected based on the recipient's vertical and horizontal corneal dimensions, which were preoperatively measured using anterior segment optical coherence tomography (Carl Zeiss Meditec, Dublin, CA, USA). The pre-cut Descemet membranes were preserved *in situ* on top of the donor corneal stroma in Optisol (Chiron, Irvine, CA, USA). The mean pre-cut-to-use period for overseas transport was 59±15 hours. Donor tissue information, including age, graft size, donor ECD, and death-to-use period is shown in Table 1.

Surgical interventions

Peripheral laser iridotomy or surgical iridectomy during cataract surgery was performed before DMEK. For patients with a cataract or a pupil anomaly, phacoemulsification with an intraocular lens (IOL) insertion or pupilloplasty was performed concomitantly or sequentially prior to DMEK without any adverse event. Endothelium-out grafts were delivered into the anterior chamber during DMEK using glass Jones tubes (GWSG, Hillsboro, OR) in 17 eyes from October 2016 to March 2019. After August 2019, a Coronet

EndoGlide insertion device (Network Medical Products, Ripon, UK) became available in Korea; after August 2019, 15 eyes received endothelium-in grafts. The size of the recipient's Descemet membrane stripping was usually 0.5–0.75 mm larger than that of the donor graft (8.0–9.0 mm).

In the endothelium-out technique, the graft was freed from the hinge and spontaneously rolled with the endothelium outward. Using a Jones tube, the graft was aspirated inside the tube with a balanced salt solution (Bausch and Lomb, Inc., Rochester, NY) and injected into the anterior chamber through a 3.5-mm corneal or scleral tunnel. The graft was unfolded using tapping procedures and was attached using an air tamponade (80–95%) after confirming that the endothelium was positioned downward by the S mark (<u>Fig</u> <u>2A</u>) [28].



Fig 2. Representative photographs of surgical procedures in Descemet membrane endothelial keratoplasty (DMEK). Endothelium-out delivery with John's tube (A) and endothelium-in technique using the DMEK EndoGlide (B). <u>https://doi.org/10.1371/journal.pone.0270037.g002</u>

In the endothelium-in technique, the graft was harvested and tri-folded inward with its endothelium. The graft was pulled into the coronary EndoGlide, which had been filled with the balanced salt solution. The graft was delivered into the anterior chamber in the correct orientation with the endothelium facing downward through a 2.75-mm corneal or scleral tunnel. The graft was unfolded with the assistance of tapping and was attached using an air tamponade (80–95%) with a confirmation of the S mark (<u>Fig 2B</u>) [14]. The standard follow-up consisted of examinations at 1 day, 2 days, 1 week, 1, 3, and 6 months, and every 6 months after DMEK.

All the patients received systemic steroid treatment for more than 3 weeks. Topical 1% prednisolone acetate (Pred Forte; Allergan, Inc., Irvine, CA, USA) was administered 4 times a day and the dose was tapered down to 1/day after 12 months. Topical 0.5% moxifloxacin eye drops (Vigamox, Alcon, TX) were applied 4 times a day.

Clinical outcome measurements

The following data were collected as the main outcomes and analyzed in each time point in a whole group and were also compared in between the endothelium-in and endothelium-out deliveries; graft survival, postoperative best corrected visual acuity (BCVA), ECD, CCT, re-bubbling rate and annual EC loss (ECL) rate. Early and delayed complications were additionally compared between those two groups. Visual acuities measured initially by Snellen charts were converted to logarithm of the minimum angle of resolution (logMAR) values for analysis. The proportions of BCVA ≥20/40, 20/25, or 20/20 were evaluated. Patients with amblyopia, advanced glaucoma, or macular disease that could affect visual acuity were excluded from visual acuity analysis. Increased IOP was defined as IOP > 21 mmHg. ECD was checked by specular microscopy (Konan Medical Inc., Hyogo, Japan). CCT was measured with anterior segment optical coherence tomography. The postoperative immediate and delayed complications were examined by slit-lamp microscopy (Haag-Streit AG, Koniz, Switzerland). To analyze the changes of corneal astigmatism, axial length (AXL), and mean numerical refractive errors (MNEs) in DMEK combined with cataract surgery (n = 18), an auto-keratorefractometer (Atlas, Carl Zeiss Meditec, Dublin, CA), topography (Orbscan II Bausch & Lomb, Claremont, CA, USA), and IOL master (Carl Zeiss MeditecAG, Jena, Germany) were used. MNEs were calculated by subtracting the value of postoperatively measured refractive error from the value of the predicted refractive error calculated by IOL master (MNE = refractive errorpredicted-refractive error_{postoperatively measured}) [21]. The correlation of either ΔCCT_{pre-post} (CCT_{preoperative}-CCT_{postoperative}) or $\Delta K_{pre-post}$ ($K_{preoperative}$ - $K_{postoperative}$) with MNE was analyzed. In most patients, the preoperative posterior curvature (K_{post}) data were not available. Therefore, postoperative Kpost (n = 15) was compared with the Kpost in eyes of normal controls (n = 50) in whom the parameters for cataract surgery were measured using an IOL Master 700 (Carl Zeiss Meditec AG, Jena, Germany).

Graft failure was defined as irreversible loss of corneal graft clarity despite intense medical treatment. Immunologic rejection of corneal grafts was diagnosed as the sudden onset of corneal edema developed in the presence of ocular inflammation (ciliary injection, endothelial rejection line, keratic precipitates, stromal infiltrates or cells in the anterior chamber).

Statistical analyses

Data are presented as mean ± standard deviation where applicable. Statistical analysis was performed using SPSS Statistics 20.0 (IBM, Armonk, NY). The chi-squared test or Fisher's exact test was used to compare the two categorical variables, and Student's ttest was applied to compare the means of continuous variables between two groups. The Kaplan-Meier curves were used to analyze the corneal graft survival after a successful DMEK and a statistical difference in Kaplan-Meier curves was assessed by Log-rank test. Pearson's bivariate co-relation was used between those ocular biometric parameters. A P value of < 0.05 was considered statistically significant.

Results

Baseline characteristics of DMEK patients

The baseline characteristics of the donors and recipients are shown in <u>Table 1</u>. The follow-up duration was significantly longer in the endothelium-out group compared to the endothelium-in group (42.3±11.8 months vs 23.0±7.1 months, p<0.001). There were no significant differences between those two groups in the recipients' age, sex ratio, preoperative BCVA, anterior chamber depth, AXL, or the proportion of preoperative diagnoses with FED or BK. Combined ophthalmic diseases were present in 20 out of 32 eyes (62.5%), and the most common disease was glaucoma (n = 11, 34.4%). The other concomitant morbidities were uveitis, toxic anterior segment syndrome, iris-fixated phakic IOL-induced iritis, cystoid macular edema, epiretinal membrane, pseudoexfoliation syndrome with zonular weakness, plateau iris, and Sjögren syndrome.

Among the donors' clinical features, the graft size was significantly larger in the endothelium-in group (8.0 ± 0.4 mm) than in the endothelium-out group (7.4 ± 0.3 mm, p<0.001). The mean donor death-to-use period in the endothelium-in group (135 ± 8 hours) was similar to that in the endothelium-out group (127 ± 12 hours, p = 0.051). The mean pre-cut-to-use period did not differ between the two groups (59 ± 15 vs. 58 ± 15 hours, respectively; p>0.05). There was no difference in the ECD between the two groups.

Overall clinical outcomes of DMEK in eyes of Asian patients.

The overall graft survival rate was 71.9% at the last follow-up (Fig <u>3A</u>). Primary graft failure occurred in only 2 patients (6.3%); rebubbles were performed in 11 eyes (n = 11, 34.4%). Overall graft failures during the follow-up were caused by multiple factors, which included primary graft failure (n = 2, 6.3%), uncontrolled immunological rejection (n = 1, 3.1%), persistent uveitis with progressive peripheral anterior synechia (n = 1, 3.1%), implantation of a shunt device (n = 3, 9.4%), and corneal bacterial infection (n = 2, 6.3%). Due to corneal edema, 75% of the recipients' preoperative ECDs could not be measured (n = 24); the measurable ECD averaged 488±79 cells/mm² (n = 8). After DMEK, the mean ECD was 1400±480 cells/mm² at 3 months, 1291±497 cells/mm² at 6 months, 1086±384 cells/mm² at 1 year, and 951±252 cells/mm² at 2 years (Fig <u>3B</u>). The postoperative BCVA significantly increased compared to the preoperative BCVA by 89.3% (0.92±0.40 vs. 0.28±0.35, respectively; p<0.001) (Fig <u>3C</u>, Table <u>2</u>). Among the postoperative BCVA tests, 75% of the total eyes showed a BCVA better than 20/40, 35.7% had a BCVA better than 20/25, and 28.6% had a BCVA better than 20/20 (Fig <u>4A</u>). The mean preoperative CCT (677±63 µm) decreased to 531±58 µm at 3 months, 550±88 µm at 6 months, 561±95 µm at 1 year, and 561±119µm at 2 years (Fig <u>3D</u>). The annual ECL rate and number of glaucoma medications are described in Table <u>2</u>. The ECL rate from the donor ECD was 47.0% at 1 month, which decreased to 28.4% by 1 year, and further decreased to 17.8% by 2 years. The cumulative EC survival rate from the ECD of the donor pre-cut tissues was shown in Fig <u>5A</u>.



Fig 3. Clinical outcome of Descemet membrane endothelial keratoplasty (DMEK) using overseas transport pre-cut tissues within 3 days. Kaplan-Meier plots of overall imported graft survival rate (A), endothelial cell density (ECD) (B), best corrected visual acuity (BCVA) (C), and central corneal thickness (CCT) (D) in total cases. Data are expressed as the mean ± standard error of the mean (B, C, D).

https://doi.org/10.1371/journal.pone.0270037.g003



Fig 4.

The proportion of eyes that achieved different levels of best spectacle-corrected visual acuity (BCVA) in Descemet membrane endothelial keratoplasty (DMEK) in all cases (A) and in between endothelium-out and endothelium-in groups (B). Percentages of eyes that achieved BCVA \geq 20/40, \geq 20/25 or \geq 20/20 were not different between those two groups. <u>https://doi.org/10.1371/journal.pone.0270037.g004</u>



Fig 5. Cumulative endothelial cell survival rate of Descemet membrane endothelial keratoplasty (DMEK) using overseas transport pre-cut tissues within 3-day pre-cut-to-use period.

Cumulative endothelial cell (EC) survival rate in total cases (A). Comparison of the cumulative EC survival rate after DMEK between endothelium-out (endo-out) and endothelium-in (endo-in) groups (B). Total 17 eyes (endo-out; n = 9 vs endo-in: n = 8) were included in 1.5-year FU and 14 eyes (endo-out; n = 9 vs endo-in: n = 5) were included in 2-year FU. There were no differences between those two groups over time. Data are expressed as the mean ± standard error of the mean (A, B). https://doi.org/10.1371/journal.pone.0270037.g005

	Total (n = 32)	Endo-out (n = 17)	Endo-in (n = 15)	P value
Immunologic rejection (n,%)	4 (12.5)	3-(17.6)	1 (6.7)	0.683
Primary graft failure (n.%)	2 (6.3)	2(11.8)	0.00	0.496
Re-bubble (n/h)	11 (34.4)	5-(29.4)	6(40.0)	0.529
Postoperative BCVA (logMAR)				
Improved VA (s,%)	25 (88.3)	11(78.6	14 (308.6)	0.222
Best BCVA during FU	0.28±0.4	0.28±0.4	0.28±0.3	0.987
No. of glasscoma drops				
Postoperative 1 yr	0.7±0.9	0.7+0.9	0.610.8	0.738
Annual ECL rate" (%)				
im"	47.0±18	90.2121	43.4±13	0.390
1 97	28.4x18	29.8+20	27.7+16	0.859
2 44	17.8±13	19.6±11	34.6±17	0.522

n, member, DL, endothenar of Idsi.org/10.1371/journal.pone.02

Table 2. Postoperative clinical outcomes. https://doi.org/10.1371/journal.pone.0270037.t002

Comparative efficacies of DMEK outcome between the endothelium-out and endothelium-in deliveries

The overall survival rate was 66.7% in the endothelium-out group and 86.7% in the endothelium-in group (Fig 6A). The mean survival times were 32.2 ± 19.7 and 21.0 ± 7.9 months in the endothelium-out and endothelium-in groups, respectively, without a significant difference (p = 0.219, Fig 6A). The EC survival rate, ECD and CCT in the endothelium-in group at each time point were comparable to those in the endothelium-out group (p>0.05, Figs <u>5B</u>, <u>6B and 6D</u>). The postoperative BCVA was 0.28±0.4 in the endothelium-out group and 0.28±0.3 in the endothelium-in group (p = 0.987, Fig 6C). The percentage of eyes with a BCVA better than 20/40 was 78.6% in the endothelium-out group and 71.4% in the endothelium-in group (p>0.05). The proportion of eyes with BCVA better than 20/25 was 42.9% and 28.6% in the endothelium-out and endothelium-in groups, respectively (p>0.05). The fractions of eyes with BCVA better than 20/20 were 28.6% and 28.6% in the endothelium-out and endothelium-out and endothelium-in groups, respectively (p>0.05, Fig 4B).

Fig 6. Comparison of the clinical outcome of descemet membrane endothelial keratoplasty (DMEK) between endothelium-out (endo-out) and endothelium-in (endo-in) groups.

Kaplan-Meier plots of graft survival rate (A), endothelial cell density (ECD) (B), best corrected visual acuity (BCVA) (C), and central corneal thickness (CCT) (D). There were no differences between those two groups over time. Data are expressed as the mean ± standard error of the mean (B, C, D).

As shown in <u>Table 2</u>, the proportions of immunological rejection, primary graft failure, and re-bubble rate did not differ significantly between the endothelium-out and endothelium-in groups (p>0.05). The annual ECL rate also did not differ between the two groups (p>0.05).

Postoperative complications after DMEK in endothelium-out and endothelium-in deliveries

The immediate and delayed postoperative complications following DMEK are presented in <u>Table 3</u>. A transiently increased IOP was the most common immediate postoperative adverse event (21.9%) and was controlled with anti-glaucoma medication within a day. Secondary glaucoma was the most common delayed complication (21.9%) (<u>Table 3</u>), and 3 patients had a glaucoma drainage device implanted for uncontrolled glaucoma. Due to intraoperative iris bleeding, one patient had anterior chamber hyphema-inducing inferior partial graft detachment in the endothelium-in group; however, spontaneous resolution was observed. Fibrin formation by intensive inflammation was observed in 2 eyes of the endothelium-out group and was successfully managed using topical and oral corticosteroids. Progressive peripheral anterior synechiae developed after DMEK in a patient with previous toxic anterior segment syndrome after cataract surgery. In the endothelium-out group, 2 patients developed corneal infectious ulcers at 6 months, which were resolved with anti-bacterial treatment. The complication rates did not differ between the endothelium-in and endothelium-out groups (p>0.05, <u>Table 3</u>).

 Table 3. Postoperative complication after DMEK.

 https://doi.org/10.1371/journal.pone.0270037.t003

Refractive status and correlation of biometric factors with refractive errors

At postoperative 3 months, the mean MNEs were +0.42±1.4 D in total (Barrett Universal II), and +0.40±1.5 D and +0.45±1.4 D in the endothelium-out and endothelium-in groups, respectively (<u>Fig 7A</u>). Postoperative keratometric astigmatism (1.3±1.6 D) was significantly reduced following DMEK (preoperative: 2.2±1.2 D, p = 0.010; <u>Fig 7B</u>). $\Delta K_{pre-post}$ was positively correlated with K_{preoperative} (r = 0.5893, p = 0.002; <u>Fig 7C</u>), suggesting that the steeper the preoperative corneal anterior curvature, the larger anterior curvature changes following DMEK. Neither $\Delta K_{pre-post}$ nor $\Delta CCT_{pre-post}$ was correlated with the MNE calculated using the Barrett Universal II formula (p>0.05; <u>Fig 7D and 7E</u>). MNE also did not correlate with the ΔAXL (p>0.05; <u>Fig 7F</u>). Mean value of the K_{post} in the eyes following DMEK was 0.01±0.15, and it was not different from the K_{post} in normal controls (+0.01±0.1D, *p* = 0.941, <u>Fig 7G</u>).



Fig 7. Change in the refractive parameters following Descemet Membrane Endothelial Keratoplasty (DMEK).

The mean numerical refractive error (MNE) was calculated by subtracting achieved postoperative spherical equivalents from attempted predicted refraction (MNE = refraction_{predicted}-refraction_{postoperative measured}). mean MNEs using Barrett Universal II formula were +0.42±1.4D in total group, +0.40±1.5D in endothelium-out group and +0.45±1.4D in endothelium-in group (A). Postoperative keratometric astigmatism was significantly reduced after DMEK (p = 0.010, B). Scatterplot presents positive correlations of preoperative corneal power (K) with $\Delta K_{pre-post}$ ($K_{preoperative}-K_{postoperative}$) measured by keratometry (p = 0.002, C). Scatterplot shows MNE was not correlated with $\Delta K_{pre-post}$, $\Delta CCT_{pre-post}$ and ΔAXL (axial length preoperative-axial length postoperative) (p>0.05, E). Graphs compares the changes of posterior corneal curvature (ΔK_{post}) in eyes with DMEK to $\Delta K_{total-ant}$ in normal eyes, which were measured by IOLMaster 700 (G). Data are expressed as the mean ± standard error of the mean (A, B, G).

https://doi.org/10.1371/journal.pone.0270037.g007

Discussion

This study demonstrates that 1) the clinical efficacy and safety of DMEK in eyes of Asian patients using imported pre-cut tissues with a < 3-day pre-cut-to-use time are comparable to those in penetrating keratoplasty for BK, 2) the efficacy of the DMEK endothelium-in method is comparable to that of the endothelium-out method, and 3) a hyperopic shift by +0.42 D and a significant reduction of corneal astigmatism were observed. Corneal edema-induced anterior curvature changes may not affect the refractive error. However, the ECL within a month and the early annual rates of ECL were greater than those previously reported [<u>17</u>, <u>29–31</u>].

Since endothelial keratoplasty, including DMEK, DSAEK, and ultrathin-DSAEK, results in better visual acuity and lower graft rejection than those by penetrating keratoplasty [2, 3, 7], endothelial keratoplasty is currently preferred for both BK and FED despite the challenging surgical technique. For DMEK in Western eyes, the 1- and 5-year survivals of grafts are reported to be 94% and 83% [3, 32, 33]. The central ECD of the graft decreased by 35–38% at 6 months and by 48%–59% at 5 years in DMEK for Western eyes [32, 34]. The annual loss rate of ECD in the graft following 1 year was -8% to -10% [32]. The proportion of visual acuity better than or equal to 20/40 has been reported as 88% and 96% in reports with more cases of FED [32, 33]. In addition to the DMEK procedures being more challenging, ECL may be greater in Asian eyes than in Western eyes due to their tight small fissures, shallow anterior chamber, high vitreous pressure, thicker pigmented iris, and the effortless formation of fibrin and posterior synechiae thereafter [8]. Considering the donor shortage and rarity of FED in Asia [10, 11, 35], a lack of experience due to the low surgical volume could also be related to the surgeon's technical difficulties and be a barrier in performing DMEK.

Initial ECL at 6 months varies from 21 to 56% in reports, based on the proportion of the included FEDs and the various techniques of graft delivery to the anterior chamber [8, 9, 16, 36, 37]. In these studies, either a donor graft was manually prepared in the operating room directly prior to surgery without any intervening period [9, 16, 36, 37], or a pre-stripped donor was provided by the Eye Bank without information on the pre-cut-to-use period [8]. In the present study, the ECD at 1 month decreased by 47% from the donor ECD before the pre-cut procedure, which is a larger reduction compared to those in previous studies [15, 36]. It may be caused by the long pre-cut-to-use time (> 48 hours), the anatomical challenges in Korean eye, or the effect of the initial learning-curve. Moreover, relatively higher ECL rate and lower survival than those of previous Western studies [15, 32, 34] may have been affected by the high proportion of combined ocular morbidities (62.5%) including glaucoma (34%) and uveitis (18.8%).

Advanced FED with stromal edema was included in this study, whereas FED eyes without significant corneal edema were included in reports from Western countries [24, 25, 34]. In this study, the ECL over time was not different between the BK and grade-4 FED (FED vs BK; 44.8±20% vs 49.0±18% at 1month; 28.1±18% vs 26.9±18% at 1st year; 16.5±11% vs 17.8±12% at 2nd year, p > 0.05). A high proportion of BK inclusions (62.5%) also contributed to the high ECL at the early stage. However, similar to a DSAEK study using imported corneas [38], this study also presented a comparable survival rate with restoration of good visual acuity, suggesting that DMEK within a 3-day pre-cut-to-use period might work well in eyes of Asian patients. Since the survival rates of penetrating keratoplasty for BK were 89.8% for the first year and 47.2% for the fifth year [39], the survival of DMEK in this study showed comparable survival to that in PKP for BK.

Currently, graft delivery and unfolding procedures need to be standardized to improve the surgical outcomes. Critical steps are still performed in variable ways among different surgeons. To minimize either delivery-associated or tapping-induced ECL, we compared the surgical outcomes between the endothelium-in and endothelium-out deliveries. The visual acuity, ECL, and CCT did not differ between the two methods. Recently, in vitro studies discovered that ECL with the tri-folded endothelium-in method was comparable to that with the scroll-based endothelium-out method [<u>40–42]</u>. Moreover, recent data from the Busin group using trifolded endothelium-in pull-through delivery demonstrated sustained survival [15]. Since the folding of the graft with the endothelium in method reduces frictional damage to the wall of the insertion cartridge, plastic cartridges, such as the IOL cartridge or EndoGlide can be used. Glass injectors, such as the Geuder cannulas or modified Jones tubes, are used for scrolled grafts with the endothelium-out method. Clinical studies have also presented acceptable ECL rates in DMEK with the endothelium-in delivery [15, 16, 43, 44]. In Korea, Geuder cannula is not commercially available. At first, we used modified Jones for a delivery of endotheliumout graft through the 3.5 mm wound. During the inserting procedure, anterior chamber tends to be easily collapsed due to larger wound, which leads to disperse iris pigments and makes unfolding procedure difficult in small-dimension eyes. After EndoGlide is commercially available in Korea, we used EndoGlide to deliver endothelium-in graft through the 2.75 mm wound and found that anterior chamber appears to be more stable with EndoGlide compared with modified Jones tube in small dimension eyes. Therefore, we aimed to compare the in vivo outcome in between endothelium-out and endothelium-in with different delivery method. This study supports the previous study by Price et al that showed no differences between the endothelium-in and endothelium-out delivery [17]

Since the dark pigmented iris blocks visibility to the graft, graft positioning with an appropriate orientation is difficult in eyes of Asian patients. Therefore, the tri-folded endothelium-in method with the assistance of graft-grabbing forceps may be an easier procedure with fewer upside-down placements than the endothelium-out method. In most cases, the scleral tunnel is preferred because of the small dimensions of the anterior chamber. Since the Geuder glass cannula is not currently available in Korea, a modified Jones tube with a 3.5 mm incision or an EndoGlide with a 2.75 mm incision are possible options. Using EndoGlide could be beneficial in maintaining anterior chamber stability during insertion compared to the Jones tube.

There were no differences in postoperative complications between the endothelium-in and endothelium-out delivery. One patient developed a pupillary block following re-bubble owing to discharge and early ambulation of the patient. Inferior iridectomy is usually recommended for preventing air- or gas-induced pupillary blocks [45-47]. However, peripheral iridectomy-less DMEK has also been reported without a pupillary block [48]. Since half the amount of injected air is absorbed within 12–24 hours [49, 50], peripheral iridectomy might not be necessary if the patient can maintain a supine position on the operative day for more than 6 hours. However, severe postoperative inflammation could cause fibrin formation when combined with uveitis [51]. Fibrin formation around the pupil or intraoperative bleeding can induce a pupillary block. Therefore, a peripheral iridectomy should be performed for early ambulation of the host or for the reduction of a pupillary block by air or fibrin.

This study demonstrated a hyperopic shift (+0.42 D) which corresponded with the refractive outcomes (+0.55 ~ +1.12 Diopter) of previous studies following triple DMEK [20, 22, 52, 53]. The causes of hyperopic shift remain controversial. Thicker corneal thickness [20], flatter posterior center curvature [23], preoperative posterior curvature [22], posterior-to-anterior corneal curvature radii ratio, and oblate posterior corneal profile [54] are reported to be involved in the hyperopic shift. In this study, although the positive correlation between $K_{preoperative}$ and $\Delta K_{pre-post}$ indicates anterior curvature changes between before and after DMEK, changes of neither CCT nor anterior curvature correlated with the refractive shift, which supports the study by Augustine et al [22]. Another possibility is that keratometry or axial length are measured less accurately owing to epithelial edema of BK, unlike the previous studies that included patients with FED who could maintain a regular anterior surface. However, we found no difference in the AXL between pre- and post-DMEK (S1 Fig), nor the correlation of the Δ AXL with the refractive shift. There was no difference in the posterior corneal curvature (K_{post}) between the post-DMEK and normal control eyes, suggesting that the graft does not induce a major change. However, we could not evaluate directly whether the individual changes of the posterior curvature could affect the refractive shift. Considering that previous studies show that the posterior cornea curvature correlates with the hyperopic shift of post-DMEK eyes [22, 23, 54], the posterior curvature may be a weighting factor in the refractive shift.

This study has several limitations. Owing to the limitations of its retrospective design, some of the data were missing. The small sample size owing to the low popularity of DMEK surgery could have affected the statistical power. However, considering that other studies comparing two groups with similar sample sizes [40, 42, 55], the sample size of this study seems to be adequate to render a statistically sufficient power. Due to rarity of FED cases in Asia, inevitably, the indications for surgery have diversified. Given that there are other studies that have been conducted with various surgical indications, so our study also worthy to compare the clinical outcomes [36, 56-58]. In addition, the follow-up durations differed between the two groups. Both limited and different follow-up durations may increase the risk of bias in the survival analysis. Finally, ΔK_{post} was not measured since the preoperative data were not available. Therefore, further studies to assess the correlation of the ΔK_{post} with MNE are warranted. Lastly, DMEK was performed by a professor (MK Kim) who had previous experience of performing approximately 1000 penetrating keratoplasties and 30 deep anterior lamellar keratoplasties. Before starting DMEK, the professor had undergone hands-on training programs by a DMEK wet lab course conducted by the Gerrit Melles of Netherlands Institute for Innovative Ocular Surgery (NIIOS) Academy in Rotterdam. This study reported an incidence of learning curve related-early graft failure of 6.3%, which was comparable to the previous incidences of learning curve effect, it is still noteworthy since it reports that imported pre-cut tissue is feasible for DMEK in Asian countries where the shortage of corneal donor limits its popularity.

In conclusion, performing DMEK using imported pre-cut tissue with a ≤ 3-day pre-cut-to-use period may be a feasible option for either BK or FED in eyes of Asian patients. Additionally, the tri-folded endothelium-in method seems to be comparatively effective to the endothelium-out method for DMEK with imported pre-cut tissue. A hyperopic shift may not be related to the anterior curvature changes.

Supporting information

<u>S1 Fig.</u> Graphs comparing the changes of the axial length (AXL) in eyes with Descemet membrane endothelial keratoplasty (DMEK) between preoperative (pre-op) and postoperative (post-op) groups, which were measured by IOLMaster 700. Data are expressed as mean ± standard error of the mean. <u>https://doi.org/10.1371/journal.pone.0270037.s001</u> (TIF)

References

- Melles GR, Eggink FA, Lander F, Pels E, Rietveld FJ, Beekhuis WH, et al. A surgical technique for posterior lamellar keratoplasty. Cornea. 1998;17(6):618–26. Epub 1998/11/20. pmid:9820943. <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 2. Price MO, Gupta P, Lass J, Price FW Jr. EK (DLEK, DSEK, DMEK): New Frontier in Cornea Surgery. Annu Rev Vis Sci. 2017;3:69–90. Epub 2017/07/13. pmid:28697678.

View Article • PubMed/NCBI • Google Scholar

- Price MO, Mehta JS, Jurkunas UV, Price FW Jr. Corneal endothelial dysfunction: Evolving understanding and treatment options. Prog Retin Eye Res. 2021;82:100904. Epub 2020/09/26. pmid:32977001.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 4. Melles GR, Lander F, Rietveld FJ. Transplantation of Descemet's membrane carrying viable endothelium through a small scleral incision. Cornea. 2002;21(4):415–8. Epub 2002/04/26. pmid:11973393.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 5. Melles GR, Ong TS, Ververs B, van der Wees J. Descemet membrane endothelial keratoplasty (DMEK). Cornea. 2006;25(8):987–90. Epub 2006/11/15. pmid:17102683.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 6. Melles GR. Posterior lamellar keratoplasty: DLEK to DSEK to DMEK. Cornea. 2006;25(8):879–81. Epub 2006/11/15. pmid:17102659.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 7. Stuart AJ, Romano V, Virgili G, Shortt AJ. Descemet's membrane endothelial keratoplasty (DMEK) versus Descemet's stripping automated endothelial keratoplasty (DSAEK) for corneal endothelial failure. Cochrane Database Syst Rev. 2018;6:CD012097. Epub 2018/06/26. pmid:29940078.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 8. Hayashi T, Oyakawa I, Kato N. Techniques for Learning Descemet Membrane Endothelial Keratoplasty for Eyes of Asian Patients With Shallow Anterior Chamber. Cornea. 2017;36(3):390–3. Epub 2016/12/22. pmid:28002113; PubMed Central PMCID: PMC5291281.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 9. Ang M, Ting DSJ, Kumar A, May KO, Htoon HM, Mehta JS. Descemet Membrane Endothelial Keratoplasty in Asian Eyes: Intraoperative and Postoperative Complications. Cornea. 2020;39(8):940–5. Epub 2020/05/27. pmid:32452991.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- Soh YQ, Kocaba V, Pinto M, Mehta JS. Fuchs endothelial corneal dystrophy and corneal endothelial diseases: East meets West. Eye (Lond). 2020;34(3):427–41. Epub 2019/07/04. pmid:31267087; PubMed Central PMCID: PMC7042366.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- Musch DC, Niziol LM, Stein JD, Kamyar RM, Sugar A. Prevalence of corneal dystrophies in the United States: estimates from claims data. Invest Ophthalmol Vis Sci. 2011;52(9):6959–63. Epub 2011/07/28. pmid:21791583; PubMed Central PMCID: PMC3175990. <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 12. Kim YS, Jin SY, Chung JK. Descemet Membrane Endothelial Keratoplasty for Epithelial Downgrowth After Clear Corneal Cataract Surgery. Eye Contact Lens. 2018;44 Suppl 1:S326–s9. Epub 2016/09/09. pmid:27607149.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 13. Yum HR, Kim MS, Kim EC. Retrocorneal membrane after Descemet membrane endothelial keratoplasty. Cornea. 2013;32(9):1288–90. Epub 2013/06/25. pmid:23792439.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- Busin M, Leon P, Scorcia V, Ponzin D. Contact Lens-Assisted Pull-Through Technique for Delivery of Tri-Folded (Endothelium in) DMEK Grafts Minimizes Surgical Time and Cell Loss. Ophthalmology. 2016;123(3):476–83. Epub 2015/12/22. pmid:26686969.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 15. Yu AC, Myerscough J, Spena R, Fusco F, Socea S, Furiosi L, et al. Three-Year Outcomes of Tri-Folded Endothelium-In Descemet Membrane Endothelial Keratoplasty With Pull-Through Technique. Am J Ophthalmol. 2020;219:121–31. Epub 2020/07/14. pmid:32659279.
 <u>View Article • PubMed/NCBI • Google Scholar</u>
- 16. Tan TE, Devarajan K, Seah XY, Lin SJ, Peh GSL, Cajucom-Uy HY, et al. Descemet Membrane Endothelial Keratoplasty With a Pull-Through Insertion Device: Surgical Technique, Endothelial Cell Loss, and Early Clinical Results. Cornea. 2020;39(5):558–65. Epub 2020/01/31. pmid:31996538. <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 17. Price MO, Lisek M, Kelley M, Feng MT, Price FW Jr. Endothelium-in Versus Endothelium-out Insertion With Descemet Membrane Endothelial Keratoplasty. Cornea. 2018;37(9):1098–101. Epub 2018/06/05. pmid:29863544.

View Article • PubMed/NCBI • Google Scholar

- McKee HD, Jhanji V. Learning DMEK From YouTube. Cornea. 2017;36(12):1477–9. Epub 2017/10/05. pmid:28976416. <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 19. Newman LR, DeMill DL, Zeidenweber DA, Mayko ZM, Bauer AJ, Tran KD, et al. Preloaded Descemet Membrane Endothelial Keratoplasty Donor Tissue: Surgical Technique and Early Clinical Results. Cornea. 2018;37(8):981–6. Epub 2018/06/05. pmid:29863542.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 20. Bae SS, Ching G, Holland S, McCarthy M, Ritenour R, Iovieno A, et al. Refractive Outcomes of Descemet Membrane Endothelial Keratoplasty Combined With Cataract Surgery in Fuchs Endothelial Dystrophy. J Refract Surg. 2020;36(10):661–6. Epub 2020/10/10. pmid:33034358.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 21. Karabela Y, Eliacik M, Kocabora MS, Erdur SK, Baybora H. Predicting the refractive outcome and accuracy of IOL power calculation after phacoemulsification using the SRK/T formula with ultrasound biometry in medium axial lengths. Clin Ophthalmol. 2017;11:1143–9. Epub 2017/07/04. pmid:28670106; PubMed Central PMCID: PMC5479261.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 22. Augustin VA, Weller JM, Kruse FE, Tourtas T. Can we predict the refractive outcome after triple Descemet membrane endothelial keratoplasty? Eur J Ophthalmol. 2019;29(2):165–70. Epub 2018/07/12. pmid:29991291.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 23. Fritz M, Grewing V, Böhringer D, Lapp T, Maier P, Reinhard T, et al. Avoiding Hyperopic Surprises After Descemet Membrane Endothelial Keratoplasty in Fuchs Dystrophy Eyes by Assessing Corneal Shape. Am J Ophthalmol. 2019;197:1–6. Epub 2018/09/10. pmid:30195893.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 24. Krachmer JH, Purcell JJ Jr., Young CW, Bucher KD. Corneal endothelial dystrophy. A study of 64 families. Arch Ophthalmol. 1978;96(11):2036–9. Epub 1978/11/01. pmid:309758.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 25. Repp DJ, Hodge DO, Baratz KH, McLaren JW, Patel SV. Fuchs' endothelial corneal dystrophy: subjective grading versus objective grading based on the central-to-peripheral thickness ratio. Ophthalmology. 2013;120(4):687–94. Epub 2013/02/02. pmid:23369486; PubMed Central PMCID: PMC3618600. <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 26. Boynton GE, Woodward MA. Eye-bank preparation of endothelial tissue. Curr Opin Ophthalmol. 2014;25(4):319–24. Epub 2014/05/20. pmid:24837574; PubMed Central PMCID: PMC4131679.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 27. Brissette A, Conlon R, Teichman JC, Yeung S, Ziai S, Baig K. Evaluation of a new technique for preparation of endothelial grafts for descemet membrane endothelial keratoplasty. Cornea. 2015;34(5):557–9. Epub 2015/03/10. pmid:25747161.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 28. Tenkman LR, Price FW, Price MO. Descemet membrane endothelial keratoplasty donor preparation: navigating challenges and improving efficiency. Cornea. 2014;33(3):319–25. Epub 2014/01/24. pmid:24452215.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 29. Nishino T, Kobayashi A, Yokogawa H, Mori N, Masaki T, Sugiyama K. A 10-year review of underlying diseases for endothelial keratoplasty (DSAEK/DMEK) in a tertiary referral hospital in Japan. Clin Ophthalmol. 2018;12:1359–65. pmid:30122887.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 30. Oellerich S, Ham L, Frank LE, Gorges S, Bourgonje VJA, Baydoun L, et al. Parameters Associated With Endothelial Cell Density Variability After Descemet Membrane Endothelial Keratoplasty. Am J Ophthalmol. 2020;211:22–30. Epub 2019/10/28. pmid:31647928.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 31. Ham L, van Luijk C, Dapena I, Wong TH, Birbal R, van der Wees J, et al. Endothelial cell density after descemet membrane endothelial keratoplasty: 1- to 2-year follow-up. Am J Ophthalmol. 2009;148(4):521–7. Epub 2009/06/27. pmid:19555921.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>

- 32. Vasiliauskaitė I, Oellerich S, Ham L, Dapena I, Baydoun L, van Dijk K, et al. Descemet Membrane Endothelial Keratoplasty: Ten-Year Graft Survival and Clinical Outcomes. Am J Ophthalmol. 2020;217:114–20. Epub 2020/04/14. pmid:32283096.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 33. Fajgenbaum MAP, Kopsachilis N, Hollick EJ. Descemet's membrane endothelial keratoplasty: surgical outcomes and endothelial cell count modelling from a UK centre. Eye (Lond). 2018;32(10):1629–35. Epub 2018/06/21. pmid:29921952; PubMed Central PMCID: PMC6189189.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 34. Zwingelberg SB, Büscher F, Schrittenlocher S, Rokohl AC, Loreck N, Wawer-Matos P, et al. Long-Term Outcome of Descemet Membrane Endothelial Keratoplasty in Eyes With Fuchs Endothelial Corneal Dystrophy Versus Pseudophakic Bullous Keratopathy. Cornea. 2021. Epub 2021/05/04. pmid:33935237.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 35. Kim M. Current Progress in corneal Xenotransplantation In Xenotransplantation NEW INSIGHTS 2017 1st Ed. INTECH. Editor Shuji Miyagawa 25-46
- 36. Lekhanont K, Pisitpayat P, Cheewaruangroj N, Jongkhajornpong P, Nonpassopon M, Anothaisintawee T. Outcomes of Descemet Membrane Endothelial Keratoplasty in Bangkok, Thailand. Clin Ophthalmol. 2021;15:2239–51. Epub 20210531. pmid:34103888; PubMed Central PMCID: PMC8180280.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 37. Ang M, Mehta JS, Newman SD, Han SB, Chai J, Tan D. Descemet Membrane Endothelial Keratoplasty: Preliminary Results of a Donor Insertion Pull-through Technique Using a Donor Mat Device. Am J Ophthalmol. 2016;171:27–34. Epub 2016/08/28. pmid:27565226.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 38. Lekhanont K, Vanikieti K, Nimvorapun N, Chuckpaiwong V. Outcomes of descemet stripping automated endothelial keratoplasty using imported donor corneas. BMC Ophthalmol. 2017;17(1):41. Epub 2017/04/07. pmid:28381247; PubMed Central PMCID: PMC5382393.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 39. Anshu A, Li L, Htoon HM, de Benito-Llopis L, Shuang LS, Singh MJ, et al. Long-Term Review of Penetrating Keratoplasty: A 20-Year Review in Asian Eyes. Am J Ophthalmol. 2021;224:254–66. Epub 2020/11/02. pmid:33129808.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 40. Parekh M, Ruzza A, Ferrari S, Ahmad S, Kaye S, Ponzin D, et al. Endothelium-in versus endothelium-out for Descemet membrane endothelial keratoplasty graft preparation and implantation. Acta Ophthalmol. 2017;95(2):194–8. Epub 2016/07/17. pmid:27422483.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 41. Barnes K, Chiang E, Chen C, Lohmeier J, Christy J, Chaurasia A, et al. Comparison of Tri-folded and Scroll-based Graft Viability in Preloaded Descemet Membrane Endothelial Keratoplasty. Cornea. 2019;38(3):392–6. Epub 2018/12/15. pmid:30550395.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 42. Ho J, Jung H, Banitt M. Quantitative and Qualitative Differences in Endothelial Cell Loss Between Endothelium-In Versus Endothelium-Out Loading in Descemet Membrane Endothelial Keratoplasty. Cornea. 2020;39(3):358–61. Epub 2019/11/07. pmid:31688201.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 43. Ighani M, Dzhaber D, Jain S, De Rojas JO, Eghrari AO. Techniques, Outcomes, and Complications of Preloaded, Trifolded Descemet Membrane Endothelial Keratoplasty Using the DMEK EndoGlide. Cornea. 2021;40(5):669–74. Epub 2021/01/21. pmid:33470675.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 44. Karakus S, Ighani M, Noparat P, Tofigh M, Chiang E, Barnes K, et al. Aspiration of Tri-folded, Endothelium-In Grafts for Descemet Membrane Endothelial Keratoplasty. Cornea. 2019. Epub 2019/02/01. pmid:30702471.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>
- 45. Price MO, Giebel AW, Fairchild KM, Price FW Jr., Descemet's membrane endothelial keratoplasty: prospective multicenter study of visual and refractive outcomes and endothelial survival. Ophthalmology. 2009;116(12):2361–8. Epub 2009/10/31. pmid:19875170.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 46. Naveiras M, Dirisamer M, Parker J, Ham L, van Dijk K, Dapena I, et al. Causes of glaucoma after descemet membrane endothelial keratoplasty. Am J Ophthalmol. 2012;153(5):958–66 e1. Epub 2012/01/31. pmid:22285014.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>

- 47. Dapena I, Ham L, Droutsas K, van Dijk K, Moutsouris K, Melles GR. Learning Curve in Descemet's Membrane Endothelial Keratoplasty: First Series of 135 Consecutive Cases. Ophthalmology. 2011;118(11):2147–54. Epub 2011/07/23. pmid:21777980.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 48. Livny E, Bahar I, Levy I, Mimouni M, Nahum Y. "PI-less DMEK": results of Descemet's membrane endothelial keratoplasty (DMEK) without a peripheral iridotomy. Eye (Lond). 2018. Epub 2018/12/07. pmid:30518974.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 49. Sim DA, Wong R, Griffiths MF. Injecting an air bubble at the end of sutureless cataract surgery to prevent inflow of ocular surface fluid. Eye (Lond). 2007;21(11):1444–5. Epub 2007/09/04. pmid:17767137.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 50. Ayintap E, Keskin U, Sadigov F, Coskun M, Ilhan N, Motor S, et al. The injection of air/oxygen bubble into the anterior chamber of rabbits as a treatment for hyphema in patients with sickle cell disease. J Ophthalmol. 2014;2014:696302. Epub 2014/05/09. pmid:24808955; PubMed Central PMCID: PMC3997984.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 51. Benage M, Korchak M, Boyce M, Mayko ZM, Bauer A, Straiko MD, et al. Intraoperative fibrin formation during Descemet membrane endothelial keratoplasty. Am J Ophthalmol Case Rep. 2020;18:100686. Epub 2020/04/16. pmid:32292835; PubMed Central PMCID: PMC7149398.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 52. Schoenberg ED, Price FW Jr., Miller J, McKee Y, Price MO. Refractive outcomes of Descemet membrane endothelial keratoplasty triple procedures (combined with cataract surgery). J Cataract Refract Surg. 2015;41(6):1182–9. Epub 2015/06/23. pmid:26096520.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 53. Augustin VA, Weller JM, Kruse FE, Tourtas T. Refractive Outcomes After Descemet Membrane Endothelial Keratoplasty + Cataract/Intraocular Lens Triple Procedure: A Fellow Eye Comparison. Cornea. 2021;40(7):883–7. Epub 2020/11/10. pmid:33165015.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 54. Diener R, Eter N, Alnawaiseh M. Using the posterior to anterior corneal curvature radii ratio to minimize the risk of a postoperative hyperopic shift after Descemet membrane endothelial keratoplasty. Graefes Arch Clin Exp Ophthalmol. 2020;258(5):1065–71. Epub 2020/02/02. pmid:32006090.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 55. Rickmann A, Boden K, Wahl S, Trouvain A, Schulz A, Szurman P. Comparison of clinical outcomes after precut DMEK with or without dextran-containing medium compared to standard DMEK: a prospective pilot study. Int Ophthalmol. 2022;42(2):401–9. Epub 20211021. pmid:34674086.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 56. Droutsas K, Lazaridis A, Kymionis GD, Chatzistefanou K, Moschos MM, Koutsandrea C, et al. Comparison of endothelial cell loss and complications following DMEK with the use of three different graft injectors. Eye (Lond). 2018;32(1):19–25. Epub 20171117. pmid:29148524; PubMed Central PMCID: PMC5770720.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 57. Jansen C, Zetterberg M. Descemet Membrane Endothelial Keratoplasty versus Descemet Stripping Automated Keratoplasty—Outcome of One Single Surgeon's More Than 200 Initial Consecutive Cases. Clin Ophthalmol. 2021;15:909–21. Epub 20210302. pmid:33688159; PubMed Central PMCID: PMC7936678.
 <u>View Article</u> PubMed/NCBI <u>Google Scholar</u>
- -
- 58. Romano V, Pagano L, Gadhvi KA, Coco G, Titley M, Fenech MT, et al. Clinical outcomes of pre-loaded ultra-thin DSAEK and pre-loaded DMEK. BMJ Open Ophthalmol. 2020;5(1):e000546. Epub 20201016. pmid:33094167; PubMed Central PMCID: PMC7569929.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 59. Hamzaoglu EC, Straiko MD, Mayko ZM, Sáles CS, Terry MA. The First 100 Eyes of Standardized Descemet Stripping Automated Endothelial Keratoplasty versus Standardized Descemet Membrane Endothelial Keratoplasty. Ophthalmology. 2015;122(11):2193–9. Epub 2015/08/15. pmid:26271841.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 60. Pereira NC, Gomes JÁ P, Moriyama AS, Chaves LF, Forseto ADS. Descemet Membrane Endothelial Keratoplasty Outcomes During the Initial Learning Curve of Cornea Fellows. Cornea. 2019;38(7):806–11. Epub 2019/04/30. pmid:31033696.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>

- 61. Wubbels RJ, Remeijer L, Engel A, van Rooij J. The learning curve for Descemet membrane endothelial keratoplasty performed by two experienced corneal surgeons: a consecutive series of 40 cases. Acta Ophthalmol. 2020;98(1):74–9. Epub 2019/06/05. pmid:31161732.
 <u>View Article</u> <u>PubMed/NCBI</u> <u>Google Scholar</u>
- 62. Fliotsos MJ, Campbell JA, Li X, Engstrom MB, Stoeger CG, McClory C, et al. Trends in Early Graft Failure Leading to Regrafting After Endothelial Keratoplasty in the United States. Cornea. 2021. Epub 2021/08/10. pmid:34369391.
 <u>View Article</u> • <u>PubMed/NCBI</u> • <u>Google Scholar</u>